

Salpakoski A, Portegijs E, **Kallinen M**, Sihvonen S, **Kiviranta I**, Alen M, Rantanen T, Sipilä S. Physical inactivity and pain in older men and women with hip fracture history. *Gerontology*. 2011;57(1):19-27.

Pekkanen L, Kautiainen H, Ylinen J, Salo P, Häkkinen A. Reliability and Validity Study of the Finnish Version 2.0 of the Oswestry Disability Index. *Spine* 2011;36:332-8.

Mäntyselkä P, **Korniloff K**, Saaristo T, Koponen H, Eriksson J, Puolijoki H, Timonen M, Sundvall J, **Kautiainen H, Vanhala M**. Association of depressive symptoms with impaired glucose regulation, screen-detected, and previously known type 2 diabetes: findings from the Finnish D2D Survey. *Diabetes Care*. 2011 Jan;34(1):71-6.

Karavirta L, Häkkinen K, Kauhanen A, Arijä-Blázquez A, Sillanpää E, Rinkinen N, **Häkkinen** Individual responses to combined endurance and strength training in older adults. *Med Sci Sports Exerc*. 2011 Mar;43(3):484-90

Sallinen J, Mänty M, Leinonen R, Kallinen M, Törmäkangas T, Heikkinen E, Rantanen T. Factors associated with maximal walking speed among older community-living adults. *Aging Clin Exp Res*. 2011 Aug;23(4):273-8.

Uutela T, Hannonen P, Kautiainen H, Hakala M, Häkkinen A. Sustained improvement of health-related quality of life in patients with early rheumatoid arthritis: a ten-year follow-up study. *Clin Exp Rheumatol*. 2011 Jan-Feb;29(1):65-71.

Ylinen J. Hoitotakuu jää usein toteutumatta TULE-sairauksien kuntoutuksessa. *Suomen Lääkärilehti* 2011; 66:1201-6.

Vuorenmaa M, Häkkinen a, Paloneva J, Kiviranta I, Kautiainen H, Oikari M, Ylinen J. Preoperatiivinen fysioterapia ja olkapään pinnetilan leikkauksesta toipuminen. *Duodecim* 2011;127:935-40.

Häkkinen A, Borg H, Hakulinen M, Jurvelin J, Anttila E, **Parviainen T, Kiviranta I**. Bone mineral density of the proximal femur after hip resurfacing arthroplasty: 1-year follow-up. *BMC Musculoskeletal disorders* 2011;12:100

Neva M, **Häkkinen A**, Isomäki P, **Sokka T**. Chronic back pain in patients with rheumatoid arthritis and in a control population: prevalence and disability – a 5-year follow-up. *Rheumatology* 2011;50:1635-9.

Korniloff K, Häkkinen A, Kautiainen H, Koponen H, Peltonen M, Mäntyselkä P, Oksa H, Kampman O, **Vanhala M**. Leisure-time physical activity and metabolic syndrome plus depressive symptoms in the FIN-D2D survey. *Prev Med*. 2011 Sep 1;53(3):211.

Salo P, Ylinen J, Kautiainen H, Häkkinen K, **Häkkinen A**. Neck Muscle Strength and Mobility of the Cervical Spine as Predictors of Neck Pain: A Prospective 6 -Year Study. *Spine (Phila Pa 1976)*. 2011 Oct 21. [Epub ahead of print]

Holviala J, **Häkkinen A, Nyman K**, Aho J, Karavirta L, Häkkinen K. Load carrying walking test and its relationships to endurance and neuromuscular capabilities in women and men of different ages. *J Sports Med Phys Fitness*. 2011;51:136-44.

Karavirta L, **Häkkinen A**, Sillanpää E, García-López D, Kauhanen A, Haapasaari A, Alen M, Pakarinen A, Kraemer WJ, Izquierdo M, Gorostiaga E, Häkkinen K. Effects of combined endurance and strength training on muscle strength, power and hypertrophy in 40-67-year-old men. *Scand J Med Sci Sports*. 2011 Jun;21(3):402-11.

Sipila S, Salpakoski A, Edgren J, Heinonen A, Kauppinen MA, **Arkela-Kautiainen M**, Sihvonen SE, **Pesola M**, Rantanen T, **Kallinen M**. Promoting mobility after hip fracture (ProMo): study protocol and selected baseline results of a year-long randomized controlled trial among community-dwelling older people. *BMC Musculoskelet Disord*. 2011 Dec 7;12(1):277.

Salo P. Mittaaminen niskapotilaan hoidossa. *Fysioterapia* 2011;58(2):4-8.
Tarnanen S, Neva M, **Kautiainen H, Ylinen J, Pekkanen I**, Kaistila L, **Vuorenmaa M, Häkkinen A**: Lannerangan jäykistysleikkauksen vaikutus selän toimintaan -3 kuukauden seurantatulokset. *Suomen Ortopedia ja Traumatologia* 2011;34:157-159.

Pekkanen L, Neva M, **Kautiainen H, Kyrölä K**, Marttinen I, Wahlman M, Vihtonen K, **Häkkinen A**. Lannerangan jäykistysleikkattujen toimintakyky ja elämänlaatu: *Suomen Ortopedia ja Traumatologia*, 2011;34:160-162.

Vuorenmaa M, Kallinen M, Ylinen J, Kautiainen H, Heinonen A, **Häkkinen A.** Mini-interventiolla ei vaikutusta 2-tyypin diabeteksen riskihenkilöiden liikunta-aktiivisuuteen tai elämänlaatuun. *Fysioterapia* 2011;58 (1):27-31.

Piirilä P, Laiho MK, **Mustonen P,** Graner M, Piilonen A, Harjola V-P, Sovijärvi ARA. The membrane component of diffusing capacity is reduced in pulmonary embolism – association with the extent of embolism. *Clin Physiol Funct Imaging* 2011; 31:196-202.

Rimpiläinen R, Hautala N, Koskenkari J, Mosorin M, Rimpiläinen J, Ohtonen PP, **Mustonen P,** Surcel H-M, Savolainen E-R, Ala-Kokko TI, Juvonen T. Minimized cardiopulmonary bypass reduces microembolization determined with retinal fluorescein angiography. *Ann Thorac Surg* 2011;91:16-22.

Korkeila P, **Mustonen P.** Antikoagulaatio ja verihutaleiden estäjät tahdistinasennuksen yhteydessä. *Sydänääni* 2011;22:23-34.

Syvänne M, Airaksinen J, Halinen M, Kosonen P, Kuusisto J, Laine M, Lammintausta O, Lepäntalo A, **Mustonen P,** Niemelä M, Pietilä M, Tierala I, Vikman S. Sepelvaltimotaudin antitromboottinen hoito, *Lääkärilehti* 2011(8) 635-42.

Rimpiläinen R, Hautala N, Koskenkari JK, Rimpiläinen J, Ohtonen PP, **Mustonen P,** Surcel HM, Savolainen ER, Mosorin M, Ala-Kokko TI, Juvonen T. Comparison of use of minimized cardiopulmonary bypass with conventional techniques on the incidence of retinal microemboli during aortic valve replacement surgery. *Perfusion.* 2011;26:479-86.

Piira OP, **Mustonen PE,** Miettinen JA, Huikuri HV, Tulppo MP. Leisure time emotional excitement increases endothelin-1 and interleukin-6 in cardiac patients. *Scand Cardiovasc J* 2012 Feb;46(1):7-15.

Burn J, Gerdes AM, Macrae F, **Mecklin JP,** Moeslein G, Olschwang S, Eccles D, Evans DG, Maher ER, Bertario L, Bisgaard ML, Dunlop MG, Ho JW, Hodgson SV, Lindblom A, Lubinski J, Morrison PJ, Murday V, Ramesar R, Side L, Scott RJ, Thomas HJ, Vasen HF, Barker G, Crawford G, Elliott F, Movahedi M, **Pylvanainen K,** Wijnen JT, Fodde R, Lynch HT, Mathers JC, Bishop DT; CAPP2 Investigators.

Long-term effect of aspirin on cancer risk in carriers of hereditary colorectal cancer: an analysis from the CAPP2 randomised controlled trial. *Lancet*. 2011 Dec 17;378(9809):2081-7.

Nieminen TT, Abdel-Rahman WM, Ristimäki A, Lappalainen M, Lahermo P, **Mecklin JP**, Järvinen HJ, Peltomäki P. BMPR1A mutations in hereditary nonpolyposis colorectal cancer without mismatch repair deficiency. *Gastroenterology*. 2011 Jul;141(1):e23-6.

Turtiainen J, Saimanen EIT, Partio TJ, Mäkinen KT, Reinikainen MT, **Virkkunen JJ**, Vuorio KS, Hakala T. Supplemental postoperative oxygen in the prevention of surgical wound infection after lower limb vascular surgery: a randomized controlled trial. *World J Surg* 2011;35:1387-1395

Niittymäki I, Tuupanen S, Li Y, Järvinen H, **Mecklin JP**, Tomlinson IP, Houlston RS, Karhu A, Aaltonen LA. Systematic search for enhancer elements and somatic allelic imbalance at seven low-penetrance colorectal cancer predisposition loci. *BMC Med Genet*. 2011 Feb 14;12:23.

Niittymäki I, Gylfe A, Laine L, Laakso M, Lehtonen HJ, Kondelin J, Tolvanen J, Nousiainen K, Pouwels J, Järvinen H, Nuorva K, **Mecklin JP**, Mäkinen M, Ristimäki A, Ørntoft TF, Hautaniemi S, Karhu A, Kallio MJ, Aaltonen LA. High frequency of TTK mutations in microsatellite-unstable colorectal cancer and evaluation of their effect on spindle assembly checkpoint. *Carcinogenesis*. 2011 Mar;32(3):305-11.

Aktan-Collan KI, Kääriäinen HA, Koltola EM, **Pylvänäinen K**, Järvinen HJ, Haukkala AH, **Mecklin JP**. Sharing genetic risk with next generation: mutation-positive parents' communication with their offspring in Lynch Syndrome. *Fam Cancer*. 2011 Mar;10(1):43-50.

Okkonen M, Varpula M, Linko R, Perttilä J, Varpula T, Penttilä V and the FINNALI study group* appendix: **Raili Laru-Sompa**. N-terminal-pro-BNP in critically ill patients with acute respiratory failure: a prospective cohort study. *Acta Anaesthesiol Scand* 2011;1-9.

Romanov M. Onko lapsi heitetty pesuveden mukana? *Suomen Lääkärilehti* 2011;66;342-343.

Forslund T, Kellokumpu I, Elomaa E, Arola J, Nuorva K. Remission of membranous glomerulonephritis after pancreatectomy for pancreatic neuroendocrine neoplasm – a rare coincidence. *Clinical Nephrology* 2011;1:42-46.

Xu L, Wang Q, Wang Q, **Lyytikäinen A**, Mikkola T, Völgyi E, Cheng S, Wiklund P, Munukka E, Nicholson P, Alén M, Cheng S. Concerted actions of insulin-like growth factor 1, testosterone, and estradiol on peripubertal bone growth: A 7-year longitudinal study. *Journal of Bone and Mineral Research* 2011;26:2204-2211.

Wiklund P, Leiting X, **Lyytikäinen A**, **Saltevo J**, Wang Q, Vögyi E, Munukka S, Alen M, Keinänen-Kiukaanniemi S, Cheng S. Prolonged breast-feeding protects mothers from later-life obesity and related cardio-metabolic disorders. *Public Health Nutrition* 2011;23:1-8.

Rintala M, **Lyytikäinen A**, Leskinen T, Alen M, Pietiläinen KH, Kaprio J, Kujala UM. Leisure-time physical activity and nutrition: a twin study. *Public Health Nutr* 2011;14;846-52

Völgyi E, Alen M, Xu L, **Lyytikäinen A**, Wang Q, Munukka E, Wiklund P, Tylavsky FA, Cheng S. Effect of long-term leisure time physical activity on lean mass and fat mass in girls during adolescence. *J Appl Physiol* 2011;110:1211-8.

Korpi-Hyövälti E, Schwab U, Laaksonen DE, Linjama H, Heinonen S, **Niskanen L**. Effect of intensive counselling on the quality of dietary fats in pregnant women at high risk of gestational diabetes mellitus. *Br J Nutr*. 2011 Nov 18:1-8.

Törrönen R, Sarkkinen E, Niskanen T, Tapola N, Kilpi K, **Niskanen L**. Postprandial glucose, insulin and glucagon-like peptide 1 responses to sucrose ingested with berries in healthy subjects. *Br J Nutr*. 2011 Sep 20:1-7.

Sarkkinen ES, Kastarinen MJ, Niskanen TH, Karjalainen PH, Venäläinen TM, Udani JK, **Niskanen LK**. Feasibility and antihypertensive effect of replacing regular salt with mineral salt -rich in magnesium and potassium- in subjects with mildly elevated blood pressure. *Nutr J*. 2011 Sep 2;10:88.

Saltevo J, **Niskanen L**, **Kautiainen H**, **Teittinen J**, Oksa H, Korpi-Hyövälti E, Sundvall J, Männistö S, Peltonen M, Mäntyselkä P, **Vanhala M**. Serum calcium level is associated with metabolic syndrome in the general population: FIN-D2D study. *Eur J Endocrinol*. 2011 Sep;165(3):429-34.

Kujala UM, Jokelainen J, Oksa H, Saaristo T, Rautio N, Moilanen L, Korpi-Hyövälti E, **Saltevo J**, **Vanhala M**, **Niskanen L**, Peltonen M, Tuomilehto J, Uusitupa M, Keinänen-Kiukaanniemi S. Increase in physical activity and cardiometabolic risk profile change during lifestyle intervention in primary

healthcare: 1-year follow-up study among individuals at high risk for type 2 diabetes. *BMJ Open* 2011;19:1-9.

Astrup A, Carraro R, Finer N, Harper A, Kunesova M, Lean MEJ, **Niskanen L**, Rasmussen MF, Rissanen A, Rössner S, Savolainen MJ, Van Gaal L on behalf of the NN8022-1807 Investigators. Safety, tolerability and sustained weight loss over 2 years with the once-daily human GLP-1 analog, liraglutide. *International Journal of Obesity* 2011:1-12.

Martikainen JA, Soini EJO, Laaksonen DE, **Niskanen L**. Health economic consequences of reducing salt intake and replacing saturated fat with polyunsaturated fat in the adult Finnish population: estimates based on the FINRISK and FINDIET studies. *European Journal of Clinical Nutrition* 2011;65:1148-1155.

Saltevo JT, Kautiainen H, Niskanen L, Oksa H, Puolijoki H, Sundvall J, Keinänen-Kiukaanniemi S, Peltonen M, Tuomilehto J, Uusitupa M, Mäntyselkä P, **Vanhala MJ**. Ageing and associations of fasting plasma glucose and 2 h plasma glucose with HbA1g in apparently healthy population. FIN-D2D study. *Diabetes Res Clin Pract* 2011;93(3):344-9.

Seppälä J, **Vanhala M**, Kautiainen H, Eriksson J, Kampman O, Mäntyselkä P, Oksa H, Ovaskainen Y, Viikki M, Koponen H. Prevalence of metabolic syndrome in subjects with melancholic and non-melancholic depressive symptoms. A Finnish population-based study. *Affect Disord*. 2012 Feb;136(3):543-9.

the FIN-D2D Survey. Pajunen P, Kotronen A, Korpi-Hyövälti E, Keinänen-Kiukaanniemi S, Oksa H, Niskanen L, Saaristo T, **Saltevo JT**, Sundvall J, **Vanhala M**, Uusitupa M, Peltonen M Metabolically healthy and unhealthy obesity phenotypes in the general population. *BMC Public Health*. 2011 Oct 1;11:754.

Kuninkaanniemi H, Villberg J, **Vanhala M**, Poskiparta M. Behaviour-change interventions in primary care: influence on nutrition and on the metabolic syndrome definers. *Int J Nurs Pract*. 2011 Oct;17(5):470-7.

Puustinen PJ, Koponen H, Kautiainen H, Mäntyselkä P, **Vanhala M**. Psychological distress measured by the GHQ-12 and mortality: a prospective population-based study. *Scand J Public Health*. 2011 Aug;39(6):577-81.

Rautio N, Jokelainen J, Oksa H, Saaristo T, Peltonen M, **Niskanen L**, Puolijoki H, **Vanhala M**, Uusitupa M, Keinänen-Kiukaanniemi S; FIN-D2D Study Group. Socioeconomic position and effectiveness of lifestyle intervention in prevention of type 2 diabetes: one-year follow-up of the FIN-D2D project. *Scand J Public Health*. 2011 Aug;39(6):561-70.

Salopuro TM, Saaristo T, Oksa H, Puolijoki H, **Vanhala M**, Ebeling T, **Niskanen L**, Tuomilehto J, Uusitupa M, Peltonen M. Population-level effects of the national diabetes prevention programme (FIN-D2D) on the body weight, the waist circumference, and the prevalence of obesity. BMC Public Health. 2011 May 19;11:350

Sundvall J, Leiviskä J, Laatikainen T, Peltonen M, Salomaa V, **Vanhala M**, Korpi-Hyövälti E, Lauronen J, Alfthan G. The use of fasting vs. non-fasting triglyceride concentration for estimating the prevalence of high LDL-cholesterol and metabolic syndrome in population surveys. BMC Med Res Methodol. 2011 May 10;11:63.

Erkkilä J, Punkanen M, Fachner J, Ala-Ruona E, Pönttiö I, Tervaniemi M, **Vanhala M**, Gold C. Individual music therapy for depression: randomised controlled trial. Br J Psychiatry. 2011 Aug;199(2):132-9.

Pfister R, Cairns R, Erdmann E, Schneider CA; PROactive investigators*Mauno Vanhala. Prognostic impact of electrocardiographic signs in patients with Type 2 diabetes and cardiovascular disease: results from the PROactive study. Diabet Med. 2011 Oct;28(10):1206-12.

Vanhala M, Kumpula LS, Soininen P, Kangas AJ, Ala-Korpela M, **Kautiainen H**, Mäntyselkä P, **Saltevo J**. High serum adiponectin is associated with favorable lipoprotein subclass profile in 6.4-year follow-up. Eur J Endocrinol. 2011 Apr;164(4):549-52.

Puustinen PJ, Koponen H, Kautiainen H, Mäntyselkä P, **Vanhala M**. Psychological distress predicts the development of the metabolic syndrome: a prospective population-based study. Psychosom Med. 2011 Feb-Mar;73(2):158-65.

Puustinen PJ, Koponen H, Kautiainen H, Mäntyselkä P, **Vanhala M**. Psychological distress and C-reactive protein: do health behaviours and pathophysiological factors modify the association? Eur Arch Psychiatry Clin Neurosci. 2011 Jun;261(4):277-84.

Parc Y, Mabrut J-Y, Shields C on behalf of the Mallorca Group***Mecklin JP**. Surgical management of the duodenal manifestations of familial adenomatous polyposis. British Journal of Surgery 2011;98:480-484.

Miettinen M, Mäkelä S, Haapala M, Helanterä A, Helsin H, Vanttinen T, Mustonen J. Glomerulonephritis emerging shortly after Puumala hantavirus infection: a report on 7 patients. Clinical Nephrology 2011;75:550-556.

Mustila A, Korpela M, Haapala AM, Kautiainen H, Laasonen L, Möttönen T, Leirisalo-Repo M, Ilonen J, Järvenpää S, Luukkainen R, **Hannonen P** for the FIN-RACo Study group. Anti-citrullinated peptide antibodies and the progression of radiographic joint erosions in patients with early rheumatoid

arthritis treated with the FIN-RACo combination and single disease-modifying antirheumatic drug strategies. *Clinical and Experimental Rheumatology* 2011;29:500-505.

Branco JC, Cherin P, Montagne A, Bouroubi A on behalf of the Multinational Coordinator Study Group * **Hannonen P**. Longterm therapeutic response to Milnacipran treatment for fibromyalgia. A European 1-year extension study following a 3-month study. *J Rheumatol* 2011;38:1403-12.

Havulinna J, **Leppänen OV, Järvinen LN**, Göransson H. Comparison of modified Kessler tendon suture at different levels in the human flexor digitorum profundus tendon and porcine flexors and porcine extensors: an experimental biomechanical study. *The Journal of Hand Surgery* 2011;36:670

Haraoui B, Smolen JS, Aletaha D, Breedveld FC, Burmester G, Codreanu C, Da Silva JP, de Wit M, Dougados M, Durez P, Emery P, Fonseca JE, Gibofsky A, Gomez-Reino J, Graninger W, Hamuryudan V, Jannaut Peña MJ, Kalden J, Kvien TK, Laurindo I, Martin-Mola E, Montecucco C, Santos Moreno P, Pavelka K, Poor G, Cardiel MH, Stanislawska-Biernat E, Takeuchi T, van der Heijde D; **Treat to Target Taskforce* Sokka-Isler T**. Treating Rheumatoid Arthritis to Target: multinational recommendations assessment questionnaire. *Ann Rheum Dis* 2011 Nov;70(11):1999-2002

Khan NA, **Sokka T**. Declining needs for total joint replacements for rheumatoid arthritis. *Arthritis Res Ther*. 2011 Oct 19;13(5):130

Pincus T, Castrejón I, **Sokka T**. Long-term prednisone in doses of less than 5 mg/day for treatment of rheumatoid arthritis: personal experience over 25 years. *Clin Exp Rheum*. 2011 Sep-Oct;29(5 Suppl 68):S130-8

Gossec L, Paternotte S, Aanerud GJ, Balanescu A, Boumpas DT, Carmona L, de Wit M, Dijkmans BA, Dougados M, Englbrecht M, Gogus F, Heiberg T, Hernandez C, Kirwan JR, Mola EM, Cerinic MM, Otsa K, Schett G, Scholte-Voshaar M, **Sokka T**, von Krause G, Wells GA, Kvien TK. Finalisation and validation of the rheumatoid arthritis impact of disease score, a patient-derived composite measure of impact of rheumatoid arthritis: a EULAR initiative. *Ann Rheum Dis* 2011 Jun;70(6):935-42.

Sokka T. Morning stiffness and other patient-reported outcomes of rheumatoid arthritis in clinical practice. *Scand J Rheumatol Suppl*. 2011;125:23-7.

Felson DT, Smolen JS, Wells G, Zhang B, van Tuyl LH, Funovits J, Aletaha D, Allaart CF, Bathon J, Bombardieri S, Brooks P, Brown A, Matucci-Cerinic M, Choi H, Combe B, de Wit M, Dougados M, Emery P, Furst D, Gomez-Reino J, Hawker G, Keystone E, Khanna D, Kirwan J, Kvien TK, Landewé R, Listing J, Michaud K, Martin-Mola E, Montie P, Pincus T, Richards P, Siegel JN, Simon LS, **Sokka T**, Strand V, Tugwell P, Tyndall A, van der Heijde D, Verstappen S, White B, Wolfe F, Zink A, Boers M. American College of

Rheumatology/European League Against Rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. *Arthritis Rheum.* 2011 Mar;63(3):573-86.

Felson DT, Smolen JS, Wells G, Zhang B, van Tuyl LH, Funovits J, Aletaha D, Allaart CF, Bathon J, Bombardieri S, Brooks P, Brown A, Matucci-Cerinic M, Choi H, Combe B, de Wit M, Dougados M, Emery P, Furst D, Gomez-Reino J, Hawker G, Keystone E, Khanna D, Kirwan J, Kvien TK, Landewé R, Listing J, Michaud K, Martin-Mola E, Montie P, Pincus T, Richards P, Siegel JN, Simon LS, **Sokka T**, Strand V, Tugwell P, Tyndall A, van der Heijde D, Verstappen S, White B, Wolfe F, Zink A, Boers M. American college of rheumatology/european league against rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. *Ann Rheum Dis.* 2011 Mar;70(3):404-13.

Pajunen P, Kontronen A, Korpi-Hyövälti E, Keinänen-Kiukaanniemi S, Oksa H, Niskanen L, Saaristo T, **Saltevo J**, Sundvall J, **Vanhala M**, Uusitupa M, Pletonen M. Metabolically healthy and unhealthy obesity phenotypes in the general population: the FIN-D2D Survey. *BMC Public Health* 2011;11:754.

Lithovius R, Harjutsalo V, Forsblom C, Groop P on behalf of the FinnDiane Study Group***Saltevo J**. Cumulative cost of prescription medication in outpatients with type 1 diabetes in Finland. *Diabetologia* 2011;54:496-503.

Kyrö JP, Summanen PA, Harjutsalo V, Groop PH, Forsblom C, Hietala K on behalf of the FinnDiane Study group***Saltevo J**. Decline in the cumulative incidence of severe diabetic retinopathy in patients with type 1 diabetes. *Diabetes Care* 2011;34:2005-6.

Sokka T. T2T-manifesti tähtää nivelreuman parempaan hoitoon. *Suomen Lääkärilehti* 2011;66:1472-959.

Kellokumpu I. Laparoscopic surgery in the future – where are limits? *Duodecim* 2011;127:23-34.

Forslund T, Kellokumpu I, Elomaa E, Arola J, Nuorva K. Remission of membranous glomerulonephritis after pancreatectomy for pancreatic neuroendocrine neoplasm – a rare coincidence. *Clin Nephrol* 2011;75:42-6.

Walker S, Taipale RS, **Nyman K**, Kraemer WJ, Häkkinen K. Neuromuscular and hormonal responses to constant and variable resistance loadings. *Med Sci Sports Exerc* 2011;43:26-33.

Ahtiainen J, Humi JJ, Kraemer WJ, Lehti M, **Nyman K**, Selänne H, Alen M, Pakarinen a, Komulainen J, Kovanen V, Mero AA, Häkkinen K. Heavy resistance exercise training and skeletal muscle androgen receptor expression in younger and older men. *Steroids* 2011;76:183-192.

Ahtiainen JP, Lehti M, Hulmi JJ, Kraemer WJ, Alen M, **Nyman K**, Selänne H, Pakarinen A, Komulainen J, Kovanen V, Mero AA, Häkkinen K. Recovery after heavy resistance exercise and skeletal muscle androgen receptor and insulin-like growth factor-I isoform expression in strength trained men. *Journal of Strength and Conditioning Research* 2011;25:767-

Vänttinen T, Blomqvist M, **Nyman K**, Häkkinen K. Changes in body composition, hormonal status and physical fitness in 11-, 13-, and 15-year old Finnish regional youth soccer players during a two-year follow-up. *Journal of Strength and Conditioning Research* 2011;25:3342-51.

Meretoja A, Kaste M, Roine RO, Juntunen M, Linna M, Hillbom M, Marttila R, Erilä T, **Rissanen A**, Sivenius J, Häkkinen U. Trends in treatment and outcome of stroke patients in Finland from 199 to 2007. Perfect stroke, a nationwide register study. *Annals of Medicine* 2011;43:22-30.

Meretoja A, Kaste M, Roine RO, Juntunen M, Linna M, Hillbom M, Marttila R, Erilä T, **Rissanen A**, Sivenius J, Häkkinen U. Direct costs of patients with stroke can be continuously monitored on a national level. Performance, effectiveness and costs of treatment episodes in stroke (Perfect Stroke) Database in Finland. *Stroke* 2011;42(7):2007-12.

Virkamäki A, **Saltevo J**. Finnish current care guideline for diabetes: interactive approach to improve individualised treatment. *Diabetologia* 2011;54:1264-5.

Davis TME, Ting R, Best JD, Donoghoe MW, Drury PL, Sullivan DR, Jenkins AJ, O'Connell RL, Whitting MJ, Glasziou PP, Simes RJ, Kesäniemi YA, GebSKI VJ, Scott RS, Keech AC on behalf of the FIELD study investigators * **Saltevo Juha**. Effects of fenofibrate on renal function in patients with type 2 diabetes mellitus: the Fenofibrate intervention and event lowering in diabetes (FIELD) study.

Visakorpi T, Jalkanen S, Laakso M, Lehto M, Liira H, Mäki M, Penttinen J, Rauramo I, Savolainen M, **Sokka T**, Viinikka L. Kliininen tutkimus Suomessa: tutkimuksen rakenteet. *Duodecim* 2011;127:957-9.